

Thermic continentality in Slovakia and climate changes

Jozef Vilček¹, Jaroslav Škvarenina², Jaroslav Vido², Radoslav Kandrík²

1) *Department of Geography and Applied Geoinformatics, University of Presov in Presov, Slovak Republic*

2) *Department of Natural Environment, Technical University in Zvolen, Slovak Republic*

Abstract

The influence of continents and oceans plays conceptually the key role in the climate conditions of European regions. Continentality is also an important phytogeographic factor of vegetation distribution in Slovakia. This study analysed continentality development at six meteorological stations in Slovakia during the periods 1951-2013, or 1961-2013. The results showed a slight non-significant increase of continentality index during the monitored period of 63 (53) years. Based on the results of CCM 2000 climate model we cannot expect significant changes of continentality by the end of 21st century, but the climate change will be significantly manifested by the increase of maximum and minimum air temperatures.

Key words: Thermic continentality Index, monthly temperature, climate change, Slovak Carpathian region

Introduction

Continentality of climate belongs to basic climatic characteristics of an area. It specifies the influence of the continent on climate formation. The opposite of continentality is called oceanity (maritimity), which is a set of climatic features influenced by ocean. According to the meteorological dictionary (Bednář *et al.* 1993), the most distinctive feature of continentality is large amplitude of air temperatures, which is the main characteristic of thermic continentality. On the base of other climatic elements we distinguish ombic and baric continentality.

From the point of bioclimatology, geography and ecology, continentality is an important characteristic of environmental parameters. For example, it assists us in understanding complex relationships between the plant distribution and geographic position. With the help of continentality or oceanity indices, phytogeography explains the changes in vegetation conditions from oceans to the interior of continents, gradual transition from forests to steppes and semi-deserts, as well as postglacial development of vegetation (species spreading in the Boreal, or Atlantic period) Ellenberg (1988). Klötzli (1976), Shidei (1974), Plesník (2002, 2004) presents that in comparison with ocean, land is characterised by basic humidity conditions and temperature differences caused by the distance from ocean (normal continentality), as well as by the elevation and the robustness of a mountain range (alpine continentality). Ocean air masses change from the edge to the interior of mountains. This increases continentality and its impact on vegetation to such an extent that horizontal zones are formed within mountain ranges, which is called as intra-mountain zonality. The Alps are a typical example Ellenberg (1988). From the edge of the mountains the vertical structures change from mesophilous atlantic plant communities up to extremely continental communities. Due to the alpine continentality, in the Alps we can see the ecological phenomena in a range of several tens of kilometres comparable to several thousand kilometres from the Atlantic coast up to the interior of Siberia. The impact of alpine continentality and subsequently also the intra-mountain zonality can also be observed in the Tatras of the Western Carpathians (Plesník 2002). This phenomenon is more thoroughly described in forestry and plant community literature, e.g.: Domin (1931), Fleischer (1994), Pagan (1992) Pagan and Randuška (1987), Somora (1958), Svoboda (1952).

In the conditions of Czecho-Slovakia, continentality or oceanity was examined by several authors. Hradička (1933) dealt with thermic and ombric continentality. Kveták (1974) elaborated continentality of Slovakia in a complex way using several indices. Melo (2002) in Hurbanov addressed continentality in connection with climate change. He simulated future changes of continentality

using CCCM 2000 and GISS 1998 climate models. Recently, some other studies, dealing with modelling of the climatic change, used the same characteristic to describe changes in continentality of climate in the twenty first century (Melo et al. 2013).

The aim of the presented paper is to examine to development of continentality on stations situated at different elevations during the years 1951 (1961) – 2013. The partial goal was to evaluate (un)suitability of continentality as an indicator of the ongoing climate change.

Materials and methods

The work is based on the data from the Slovak Hydrometeorological Institute (SHMI). **Table 1** presents the stations included in the analysis and their geographic characteristics. From the point of terrain we can divide the stations into three groups as follows:

- Lowlands (Michalovce, Hurbanovo)
- Valleys (Rožňava, Sliač)
- Highlands (Oravská Lesná, Skalnaté Pleso).

Continentality was calculated as a simple index of continentality (I_C) following the original definition of Supan applied by Rivas-Martinez et al. (1999):

$$I_C = (T_{max} - T_{min})$$

where I_C = Continentality Index

T_{max} = mean temperature (°C) of the warmest month

T_{min} = mean temperature (°C) of the coldest month

In the Czech and Slovak meteorological literature, continentality index is described as annual amplitude of temperature, or as an annual range of monthly mean air temperatures in °C, (difference between the maximum and minimum monthly mean air temperatures). Continentality index was also calculated for the future climate represented by GCMs scenario of CCCM 2000

(Canadian Climate Centre Model) following the works of Lapin et al. (2000), Melo (2002), Melo et al. (2013).

Table 1: Main characteristic of meteorological stations, their temperature variables (monthly mean, minimum, maximum air temperatures) and annual amplitude of temperatures as continentality index (Ic)

Station		Michalovce	Hurbanovo	Rožňava	Sliáč	Oravská Lesná	Skalnaté Pleso
Geographic factors							
Altitude (m)		112	115	289	313	780	1778
Latitude		48°45′	47°52′	48°39′	48°39′	49°22′	49°11′
Longitude		21°57′	18°12′	20°32′	19°08′	19°11′	20°14′
Landform		Lowland		Valley		Mountain	
Climatic variables							
Observed period (years)		1961-2013	1951-2013	1961-2013	1951-2013	1951-2013	1961-2013
Mean annual temperature (°C)	Mean	9.4	10.3	8.7	8.2	4.9	2.1
	σ	(0.75)	(0.8)	(0.8)	(0.7)	(0.7)	(0.8)
Minimum monthly mean temperature (°C)	Mean	-3.5	-1.9	-3.9	-4.4	-6.4	-7.2
	σ	(2.1)	(2.2)	(1.8)	(2.2)	(2.4)	(2.1)
Maximum monthly mean temperature (°C)	Mean	20.7	21,3	19.8	19.2	15.2	10.8
	σ	(1.4)	(1.3)	(1.4)	(1.4)	(1.2)	(1.3)
Continentality Index Ic – annual amplitude of temperature (°C)*	Mean						
	σ	24.2 (2.5)	23.2 (2.3)	23.7 (2.2)	23.6 (2.4)	21.6 (2.6)	18.0 (2.35)
* Ic (continentality index) is the annual range of monthly mean air temperatures in °C, (difference between the maximum and minimum monthly mean air temperatures), σ - standard deviation							

Table 2: The linear trend values ((°C)/year; (°C)/observed period) and their statistical significance levels of temperatures (monthly mean, minimum, maximum air temperatures and annual amplitude of temperatures as continentality index - Ic) for the 6 meteorological stations in Slovakia

Station		Michalovce	Hurbanovo	Rožňava	Sliach	Oravská Lesná	Skalnaté Pleso
Observed period (years)		1961-2013	1951-2013	1961-2013	1951-2013	1951-2013	1961-2013
Mean annual temperature	(°C)/year	0.0305	0.0244	0.03	0.0227	0.021	0.0284
	(°C)/observed period	1.6165	1.5372	1.59	1.4301	1.323	1.5052
	Significance*	***	***	***	***	***	***
Minimum monthly mean temperature	(°C)/year	0.0403	0.0248	0.0366	0.0088	0.0281	0.0324
	(°C)/observed period	2.1359	1.5624	1.9398	0.5544	1.7703	1.7172
	Significance*	*	NS	*	NS	+	+
Maximum monthly mean temperature	(°C)/year	0.0451	0.0373	0.0477	0.0393	0.0276	0.0453
	(°C)/observed period	2.3903	2.3499	2.5281	2.4759	1.7388	2.4009
	Significance*	***	***	***	***	**	***
Continentality Index - Ic (annual amplitude of temperature)	(°C)/year	0.0049	0.0125	0.0011	0.0305	0.0005	0.0128
	(°C)/observed period	0.2597	0.7875	0.0583	1.9215	0.0315	0.6784
	Significance*	NS	NS	NS	+	NS	NS
*Significance: + p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001; NS- not significant							

Results and Discussion

Since our goal was to evaluate the continentality over the whole varied terrain of Slovakia, we selected six meteorological stations from the network of SHMI. The elevations of stations vary from 112 to 1,778 m a.s.l. **Table 1** gives detailed information about temperature conditions of the examined stations. As we see, Hurbanovo situated in Podunajská nížina (lowland) is the warmest station, while Skalnaté Pleso in the Tatras, the highest mountains of Slovakia, is the coldest station. The amplitude of air temperature is the most important characteristic of thermic continentality calculated as the difference between the monthly mean temperatures of the warmest and the coldest months in the particular year.

Mean amplitudes were evaluated for the period from 1951 to 2013, or 1961 to 2013, depending on the length of observations at a particular station (**Table 1**). The highest amplitude of air temperature was found for Michalovce (24.2 °C), which is the lowland station situated in the Eastern Slovakia. The difference between the temperature amplitudes in Podunajská nížina (lowland) and Východoslovenská nížina (lowland) is 1°C. An interesting finding was that the amplitude of air temperature of the stations situated in valleys was also high: Rožňava (23.7 °C) and Sliach (23.6 °C). This is probably the result of their inversion positions with relatively low air temperatures in winter half-years, and high summer air temperatures. Skalnaté Pleso situated in the mountains has the lowest amplitude (18.0 °C). From the statistical point of view, the amplitude is a rather conservative parameter. The value of its standard deviation is almost equal for all stations (2.2 – 2.6). We can state that our results confirmed the opinion of Gorczyński ex Kveták (1983), etc., that continentality decreases with increasing elevation, and that from the point of thermic continentality the area of Slovakia still belongs to 3rd maritime transition zone ($I_c = 10.1$ až 25.0 °C)




Figure 1: Lowland station **Hurbanovo** (115 m a. s. l.) - yearly variation, temperature pattern and trend line of the mean annual temperature (a), maximum monthly temperature (b), minimum monthly temperature (c) and continentality index I_c – annual amplitude of temperature (d) for the period 1951-2013




Figure 2: Mountain station **Skalnáté Pleso** (1778 m a. s. l.) - yearly variation, temperature pattern and trend line of the mean annual temperature (a), maximum monthly temperature (b), minimum monthly temperature (c) and continentality index I_c – annual amplitude of temperature (d) for the period 1961-2013

Table 2 evaluates the developmental trend of temperature characteristics (mean, minimum, maximum air temperature) and of temperature amplitude, i.e. continentality index. **Figures 1 (a-d)** and **2 (a-d)** present the developmental trends of temperature characteristics from 1951 (1961) to 2013 for the lowland station of Hurbanovo (115 m a. s. l.) and the highland station of Skalnáté Pleso (1,778 m a. s. l.). Other stations are characterised in **Table 2**. All analysed characteristics of air temperature and thermic continentality have an increasing trend. **Table 2** presents the results of the Student's t-test of significance concerning the correlation coefficients of the mean annual temperature, maximum monthly temperature, minimum monthly temperature and continentality index I_c for the period 1951-2013 and the trend of linear regression as well. The highest rate of mean annual temperature increase equal to $0.0305\text{ }^{\circ}\text{C}$ per year was observed in Hurbanovo. It means that over the whole period of 63 years, mean annual temperature increased by $1.62\text{ }^{\circ}\text{C}$. The slowest rate of temperature increase ($0.0305\text{ }^{\circ}\text{C}$ per year) was found in Oravská Lesná, where the temperature increased by $1.32\text{ }^{\circ}\text{C}$ over the last 53 years. The

increasing trend of mean annual temperature was significant for all stations at 99.9 %. Minimum monthly temperature also increased, but the increase was less significant and had higher variability. Maximum monthly temperature significantly increased on all stations, mostly in Rožňava (2.52 °C in 53 years). Although the developmental trend of continentality was also increasing, the increase was slight and non-significant (in the case of Sliač station it was significant at 90.0%). The increasing trend fluctuated from 0.0005 to 0.0305 °C/year.

Tab. 3: Annual amplitude of temperature as continentality index (Ic) pre referenčné obdobie a pre klimatický scenár CCCM pre roky 2030 a 2075

Years/Scenario	Hurbanovo	Michalovce	Rožňava	Sliač	Oravská Lesná	Skalnaté Pleso
1951-1980	21.6	22.8	22.3	22.1	19.8	15.5
2030*	21.4	22.6	22.1	21.9	19.6	15.3
2075*	21.2	22.4	21.9	21.7	19.4	15.1
*Scenario CCCM						

Table 3 presents the trend of continentality until the year of 2075 according to the scenario of CCCM. It is expected that the continentality of all stations will slightly decrease. The amplitudes will decrease by 0.4 °C by the year 2075. Melo (2002) presented a similar result.

Many naturalists ask themselves a question: „**why do we not observe an increase of thermic continentality during the last period of the ongoing climate change?** “ The main cause is the increase of both maximum and minimum monthly temperatures. Since the amplitude is the difference between them, it remains without significant changes. Faster rate of maximum monthly temperature increase stimulates the increasing trend. However, it is only slight because minimum temperatures grow more slowly. Another explanation comes from the geographic definition of continentality: „continental climate is a type of climate inside the land of every continental zone affected by land features“ Činčura et al. (1985). According to this definition, the fact that the thermic continentality does not change is logical, because so far the climate change

does not change the geographic distance from the ocean. Mindřáš et al. (1996) and other works presented the ongoing changes in bio-climatological zonation. For example, in the southern lowlands of Slovakia, bio-climatic conditions suitable for a new community of a xeric forest of a warm temperate zone are gradually being formed. Similarly, the bio-climatic conditions of highlands also change. The modelled scenario of CCCM for the year 2075 assumes a complete extinction of alpine communities, and their replacement by a sub-alpine very moist forest (Mindřáš et al. 1996). It follows that the thermic continentality remains more or less constant even under the conditions of changing climate. It is the bio-climatic conditions of the vegetation zone that changes. If in the future we want to include continentality in the studies dealing with climate change or the changes of bio-climatic conditions, continentality needs to be linked to a climatic zone, vegetation zone, etc. For example, the change from a warm temperate moist forest (continentality index $I_c=23$) to a warm temperate dry forest ($I_c=23$) following the change of humidity conditions, or the change from a warm temperate moist forest ($I_c=23$) to a cool temperate moist forest ($I_c=23$) following the change in temperature according to bio-climatological classification of Holdridge (1947).

Conclusion

Continentality as well as oceanity represent an important climate characteristic of a particular area. At the same time, they are also an important factor of natural vegetation distribution, not only in the postglacial period of Holocene. Thus, it is logical that a number of climatologists, geographers, geo-botanists and foresters have dealt with continentality in Slovakia. We evaluated continentality using a simple index of continentality expressed by the amplitude of air temperature defined as the difference between the monthly mean air temperatures of the warmest and the coldest months in the year. We analysed the development of continentality during the years 1951 (1961) -2013 at six meteorological stations. We selected the stations so that they represented lowlands (Michalovce, Hurbanovo), valleys (Rořňava, Sliač) and highlands

(Oravská Lesná, Skalnaté Pleso). We found only a slight non-significant increase of continentality. While the temperature of the warmest month increased by 1.74 to 2.52 °C at all stations during 63 (53) years, the temperature of the coldest month increased by 0.55 to 2.14 °C. The continentality of the year 2075 was calculated using GCMs model of CCCM 2000 following the work of Lapin et al. (2000). The results of the climatic scenario indicate that by the end of 21st century we cannot expect significant changes in continentality, although the climate change will be closely coupled with the increase of maximum and minimum air temperatures.

References

- Bednař, J., et al. (1993): Meteorologický slovník výkladový a terminologický. Academia, Ministerstvo životního prostředí ČR, Praha, 1993, p. 147.
- Činčura, J. et al. (1983): Encyklopédia Zeme. Obzor, Bratislava, p. 287.
- Domin, K. (1931). Československé bučiny. Ministerstvo zemědělství republiky Československé. Sborn. Výzk. Ústavů Zeměděl. RČS 70: 1–87.
- Ellenberg, H. (1988). Vegetation ecology of central Europe. Cambridge University Press. 731 pp.
- Fleischer, P. (1994): Lesné rastlinné spoločenstvá. In: Vološčuk, I. (ed.): Tatranský národný park: Biosférická rezervácia. Gradus, Martin. p.149–162.
- Holdridge, L. R. (1947): Determination of World Plant Formations from Simple Climatic Data. Science 105: 367-368.
- Hruďička, B. (1933): Doba polovičních srážek a periodická amplituda ročního srážkového průběhu v Československu. Spisy vydávané přírodovědeckou fakultou Masarykovy university, Brno, p. 1-22.
- Klötzli F. (1976): Grenzen von Laubwäldern in Europa. Berichte der Deutschen Botanischen Gesellschaft 89: 371–380.
- Kveták, Š. (1983): Príspevok ku kontinentalite podnebia na Slovensku. Zborník prác SHMÚ Zv. 22, Alfa, Bratislava, p. 95-218.
- Lapin, M., Melo, M., Damborská, I., Gera, M, Faško, P., (2000): Nové scenáre klimatickej zmeny pre Slovensko na báze výstupov prepojených modelov všeobecnej cirkulácie atmosféry. In.: Národný klimatický program SR, MŽP SR, 8, p. 5-34.
- Melo M. (2002) Očakávané zmeny kontinentality podnebia v 21. storočí pre Hubanovo na základe výstupov dvoch klimatických modelov. In: Rožnovský, J., Litschmann, T. (eds.): XIV. Česko-slovenská bioklimatologická konferencia, Lednice na Moravě 2.-4. září 2002, p. 312–323
- Melo, M., Lapin, M., Kapolková, H., Pecho, J., Kružicová, A. (2013): Climate Trends in the Slovak Part of the Carpathians. In: J. Kozak et al. (eds.): The

- Carpathians: Integrating Nature and Society Towards Sustainability. Springer Berlin Heidelberg, p. 131-150.
- Mindáš, J., Lapin, M., Škvarenina, J. 1996: Klimatické zmeny a lesy Slovenska. In Národný klimatický program SR, Bratislava: MŽP SR, 5, 96 p.
- Pagan, J. (1992): Lesnícka dendrológia. Vydavateľstvo Technickej univerzity, Zvolen, 343 pp.
- Pagan, J., Randuška, D. (1987): Atlas drevín. Obzor, Bratislava, I. 360pp.
- Plesník, P. (2002). Niektoré nové aspekty diverzity biosféry. Geografický časopis 54: 115-130.
- Plesník, P. (2004). Všeobecná biogeografia. Univerzita Komenského, 428 pp
- Shidei, T. (1974): Forest vegetation zones. In: Numata, M. ed.: The flora and vegetation of Japan. Kodansha, Tokyo, Japan.p. 87-124
- Rivas-Martínez, S., Sánchez-Mata, D., Costa, M. (1999): North American boreal and western temperate forest vegetation. Itinera Geobotanica 12:5-316
- Somora, J. (1958): O rozšírení niektorých lesných drevín v skupine Lomnického štítu. Vydavateľstvo Osveta, Martin. 152 pp.
- Svoboda, P. (1952). Život lesa. Praha, Brázda, 894.

Acknowledgement

This work was accomplished as a part of the projects VEGA No.: 1/0008/13, 1/0281/11, 1/0463/14 of the Ministry of Education, Science, Research and Sport of the Slovak Republic and the Slovak Academy of Science; and the projects of the Slovak Research and Development Agency No.: APVV-0423-10, APVV-0131-11 and APVV-0303-11.

Summary

Kontinentalita ako aj oceanita predstavujú dôležitú charakteristiku klímy daného územia. Zároveň sú však aj významným faktorom pre prirodzené rozšírenie vegetácie, a to nielen v postglaciálnom období holocénu. Je preto logické že problematike kontinentality na Slovensku sa venoval celý rad klimatológov, geografov, geobotanikov a lesníkov. Zhodnotili sme kontinentalitu na pomoci jednoduchého indexu kontinentality, ktorý predstavuje amplitúdu teploty vzduchu, čiže rozdiel medzi priemernou mesačnou teplotou vzduchu najteplejšieho a najchladnejšieho mesiaca v roku. Analyzovali sme vývoj kontinentality v rokoch 1951 (1961) - 2013 a to na šiestich meteorologických staniciach. Výber staníc sme uskutočnili tak aby reprezentovali nížinné polohy

Nížiny (Michalovce, Hurbanovo), kotliny (Rožňava, Sliač) ako aj pohoria (Oravská Lesná, Skalnaté Pleso). Zistili sme síce nepatrný trend rastu kontinentality, no tento nie je štatisticky významný. Zatiaľ čo teplota najteplejšieho mesiaca na všetkých sledovaných staniciach za 63 (53) rokov vzrástla v rozsahu 1.74 až 2.52 °C a teplota najchladnejšieho mesiaca vzrástla o 0.55 až 2.14 °C/sledované obdobie hodnota indexu kontinentality sa menila len nepatrne. Pre výpočet kontinentality v roku 2075 sme použili GCMs model CCCM 2000 podľa práce Lapin *et al.* (2000). Na základe výsledkov klimatického scenára nemôžeme očakávať výraznejšie zmeny kontinentality do konca 21. storočia, no klimatická zmena bude výrazne spojená s nárastom maximálnych minimálnych teplôt vzduchu.

Contact:

ŠKVARENINA Jaroslav, prof. Dr.

Department of Natural Environment,

Technical University in Zvolen, Masarykova 24

96053 Zvolen

Slovak Republic

e-mail: skvarenina(a)tuzvo.sk